

WINTER BIRD COMMUNITIES IN AFFORESTATION: SHOULD WE SPEED UP OR SLOW DOWN ECOLOGICAL SUCCESSION?

Paul B. Hamel, Daniel J. Twedt, Timothy J. Nuttle,
Christopher A. Woodson, Fred Broerman, and Joseph M. Wahome¹

Abstract—Recent assessments of afforestation on agricultural lands in the Mississippi Alluvial Valley imply the importance of quickly developing vertical forest structure to benefit wildlife. Examining this assumption, we find that mammals and birds occur through the full successional sere as targets of proactive management and control. Different species of animals thrive in structures available at different times during succession. Thus, forest managers' choices of strategies favor species' success differentially. Early successional species, particularly those avian communities occurring during winter, have heretofore been considered only in passing. However, because they occur in areas where herbaceous plants dominate vegetation structure, these communities include species otherwise rare or absent from the landscape. Extensive afforestation in the Mississippi Alluvial Valley provides ephemeral habitat for birds that winter in herbaceous areas. To provide habitat for winter birds, managers may wish to consider maintaining large tracts in herbaceous vegetation similar to that occurring 3 to 7 years after cessation of farming activities.

INTRODUCTION

Recent assessments of afforestation of agricultural lands in the Mississippi Alluvial Valley (MAV), particularly the Delta area of Mississippi, have stressed the importance of quickly attaining the physical structure and stature of forests (Schweitzer and others 1997). Implementation of land management plans designed for neotropical migratory birds in the Lower MAV (LMAV) (Mueller and others 1995) will benefit from dependable, rapid afforestation as well. Benefits of afforestation include provision of habitat for middle and late successional birds, production of pulpwood, production of sawtimber wood products, and erosion control, as well as carbon sequestration and accumulation of soil organic matter. All these benefits are positively associated with the speed with which afforestation occurs. Rapid afforestation implies swift accumulation on the landscape of the physical structure and stature of forest. It is a means of carbon sequestration (Cannell 1999b, Chang 1999) and of land rehabilitation in Amazonia (Bauch and others 1999) and elsewhere (Harrington 1999). Fast development of vertical forest structure is implicit in the environmental (Joslin and Schoenholtz 1997) and economic (Pande and others 1999, Scholtens 1998) analyses of afforestation.

Afforestation, including rapid afforestation, is assumed to be beneficial to wildlife (Boyle 1999, Cannell 1999a, Helmer 1999, Weaver and Pelton 1994, Weaver and others 1990, Willoughby and McDonald 1999). On the other hand, certain native wildlife and grazing animals can hinder afforestation

efforts (Anderson and Katz 1993, Houston 1991, Niyaz and others 1999).

Vegetation structure is an important determinant of bird species occurrence and community composition (DeGraaf 1987, DeGraaf and others 1992, James 1971). Hamel (1992) associates birds with combinations of vegetation structure, such as trees, shrubs, and herbaceous vegetation. Afforestation yields unusual elements in secondary succession, such as tall cottonwood (*Populus L. spp.*) trees and herbaceous vegetation with little woody understory. A first hypothesis is that the bird community developing in afforestation will reflect vegetation structure.

Wintering birds use the early successional herbaceous community. During rapid afforestation, the early successional periods are shorter than during natural succession. Our second hypothesis is that early successional species may not benefit from rapid afforestation of agricultural fields as much as from natural succession.

To examine the assumptions that rapid afforestation is beneficial to wildlife and that bird species occurrence is a function of vegetation structure, we consider the presence or absence of particular bird species and their communities rather than the rate of development of ecological function. As hawks and owls prey on mammals, then succession of mammal species is, in turn, dependent on bird community dynamics.²

¹ Hamel, Research Wildlife Biologist, USDA Forest Service, Southern Research Station, Stoneville, MS; Twedt, Wildlife Biologist, U.S. Department of the Interior, Geological Survey, Vicksburg, MS; Nuttle, Graduate Research Fellow, Utah State University, Department of Biology and Ecology Center, Logan, UT; Woodson, Assistant Refuge Manager, U.S. Fish and Wildlife Service, Cruger, MS; Broerman, Wildlife Biologist, U.S. Fish and Wildlife Service, Bethel, AK; and Wahome, Associate Professor, Mississippi Valley State University, Department of Natural Sciences & Environmental Health, Itta Bena, MS.

² Wahome, Joseph. 1999. Unpublished data. On file with: Department of Natural Sciences & Environmental Health, Mississippi Valley State University, 14000 Highway 82 West, Itta Bena, MS 38941.

Citation for proceedings: Holland, Marjorie M.; Warren, Melvin L.; Stanturf, John A., eds. 2002. Proceedings of a conference on sustainability of wetlands and water resources: how well can riverine wetlands continue to support society into the 21st century? Gen. Tech. Rep. SRS-50. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 191 p.

A review of literature provides a summary of expected responses of birds and small mammals to early successional habitats in MAV landscapes. Empirical observations compare responses of early successional wildlife species to four different afforestation treatments. Our primary data set involves winter bird populations in the first few years after abandonment of agricultural lands. Nuttle (1997) investigated bird communities at afforested sites of different ages in the LMAV that were established by planting oak (*Quercus* L. spp.) seedlings. Nuttle (1997) and Nuttle and Burger (1995) summarize breeding bird communities in oak plantations, some of which were 30 years old. However, we have too few data to demonstrate what happens to species during middle successional conditions occurring 20 to 60 years after initial afforestation.

METHODS

In a comparison of bird occurrence among different methods of afforestation, we focus on total abundance, density of individuals, and species richness.

Literature Analysis

The National Council of the Paper Industry for Air and Stream Improvement, Inc. (NCASI) has published annotated bibliographies on wildlife, forestry, and habitat in the East (NCASI 1993, 1999). A keyword search yielded papers dealing with wildlife response to afforestation, reforestation, forest restoration, and old-field succession in bottomland landscapes in the Southeastern United States. However, few publications treat responses to afforestation in the MAV. Studies done in other landscapes and regions can contribute insights as similar species respond to different conditions.

Empirical Observations of Winter Bird Communities

Between late October and late April from 1996 to 2000, Broerman surveyed 8 bird species on 59 afforestation sites in 9 Mississippi counties. The sites were all former agricultural land 2 to 9 years after cessation of farming activities. Some had been planted with tree seedlings, principally Nuttall oak (*Quercus nuttallii* Palmer), green ash (*Fraxinus pennsylvanica* Marshall), willow oak (*Q. phellos* L.), or water oak (*Q. nigra* L.); some had not. Herbaceous vegetation on these sites exceeded the height of woody vegetation. The survey involved using a four-wheeler to flush birds from herb or grass-dominated vegetation. During this survey he made 273 trips to the sites. Of these, 271 trips to 57 sites took place in 7 counties.

In 1999 to 2000, Hamel conducted Winter Bird Populations Studies (WBPS) (Kolb 1965) on the treatment plots in the Sharkey large-scale restoration experiment near Anguilla, Sharkey County, MS (Sharkey Site) (Schweitzer and others 1997). Hamel and Woodson made additional observations on these plots in 1998 and 1999. As a comparison of afforestation methods, the design of the Sharkey Site was randomized complete blocks design, involving three replicates each of three afforestation treatments and a natural succession and regeneration control (NAT). The treatments were: (1) sown Nuttall oak acorns (SOW), 2,500 per hectare; (2) planted Nuttall oak seedlings (PLN), 750 per

hectare; and (3) planted cottonwood stem cuttings (NUR), 750 per hectare, followed by underplanting of Nuttall oak seedlings, 375 per hectare, 2 years later. Individual treatment plots were approximately 8 ha in extent. Initial establishment of the treatments was in spring 1995, with the underplanting in the NUR plots conducted in spring 1997.

Hamel conducted a WBPS on each of the 12 plots on the Sharkey Site. His WBPS counts were 30 minutes during the morning hours on 8 days from December 1999 to March 2000. He recorded all birds seen or heard in the vegetation of the plot, or actively foraging over the plot. He visited 1 of 12 plots in a random sequence. He summarized these results as average number of individuals encountered per species per hectare.

We test the hypothesis that early successional species may not benefit from rapid afforestation as much as from natural succession. Our null hypothesis for this test is that bird species dependent on herbaceous vegetation will be found in the faster growing cottonwood plantations as well as in other treatments that are accumulating woody structure more slowly. Our data for this test consist of the abundance and occurrence of the individual species recorded on the WBPS in the different treatment plots of the Sharkey Site.

We test the hypothesis that the bird community that develops in afforestation areas reflects the presence of separate elements of vegetation structure. We compare the occurrence of bird species among the treatments on the Sharkey Site in terms of their foraging preferences from Hamel (1992) for this test. The null hypothesis for this examination is that no association between foraging substrate and occurrence by treatment will be apparent among the birds.

We test the hypothesis that accumulating vegetation structure determines the occurrence of birds in afforestation treatments. Physical vegetation structure accumulates on the experimental treatments at a predictable sequence, measured by the height of the woody vegetation, NUR>PLN>SOW>NAT. Using the data on species richness and density of individual species among treatments from the Sharkey Site, we conduct analysis of variance (ANOVA).

Our analysis uses PROC GLM or PROC ANOVA, as a randomized complete block design with treatment or block and treatment as main effects (SAS Institute Inc. 1989). Significance was accepted at $p < 0.05$; means testing was conducted using Duncan's multiple comparison of means. For tests of the difference of abundance of individual species among treatments, we accepted significance at $p < 0.10$ experiment wide with sequential Bonferroni correction.

RESULTS AND DISCUSSION

Literature Analysis

Of 38 papers in the literature search (citations in table 1), 13 deal exclusively with single species response to habitat restoration, 6 with songbirds in general, 2 with waterfowl, 3 with general vertebrate response to reforestation, 3 with effects of streamside restoration on aquatic communities, 3 with effects of fire on habitat restoration, and 4 with the

Table 1—Literature review of wildlife responses to afforestation, reforestation, and forest restoration

Category	Citation
Single species responses to restoration	
Black bear (<i>Ursus americanus</i> Pallas)	Weaver and others 1990, Weaver and Pelton 1994
Red-cockaded woodpecker [<i>Picoides borealis</i> (Vieillot)]	Cantrell and others 1995, Conner and Rudolph 1995, Gaines and others 1995, Watson and others 1995, Wilson and others 1995
American woodcock (<i>Scolopax minor</i> Gmelin)	Sepik and Blumenstock 1993
Wild turkey (<i>Meleagris gallopavo</i> L.)	Dickson 1992
Florida scrub-jay [<i>Aphelocoma coerulescens</i> (Bosc)]	Root and others 1995; Schmalzer 1993, 1994; Schmalzer and others 1994
Beaver pond reforestation	Houston 1991
Deer browse affect on reforestation	Anderson and Katz 1993
Songbirds in general	Bielefeldt and Rosenfield 1994, Nuttle and Burger 1995, Ribbeck and Hunter 1994, Tomlinson 1977, Wesley and others 1976
Restoration for songbirds	Ribbeck and Hunter 1994
Reforestation and waterfowl	Kaminski and others 1993, Reinecke 1994
Bird dispersal of fruiting plants into restoration areas	Robinson and Handel 1993
General vertebrate response to reforestation	Askins and Philbrick 1987, Litvaitis 1993
Using vertebrates to assess cumulative impacts	Croonquist and Brooks 1991
Land management planning	McCullough 1994
Forest restoration of streamside zones, effect on aquatic species	Flebbe and Dolloff 1995, O'Brien-White and Thomason 1995, Sweeney 1993
Fire effects in restoration	Fitzgerald and Tanner 1992, Provencher and others 1995, Simberloff 1993
Old-growth forest restoration, biodiversity conservation	Harris and Scheck 1991; Mladenoff and others 1993, 1994; Vora 1994
Clearcuts	Hurst and Bourland 1996, Mitchell 1989

importance of reforestation to the maintenance of old-growth conditions on the landscape. Three papers discuss the role of animals, specifically beaver (*Castor canadensis* Kuhl), white-tailed deer [*Odocoileus virginianus* (Zimmermann)], and fruit-eating birds as agents modifying plant community composition. Predetermined objectives are essential in restoration planning, as in all other land management, a point made by a single paper (McCullough 1994). No paper

explicitly treats early secondary succession on abandoned farmland in the MAV. The greater proportion of the papers indicate that restoration that proceeds faster from open ground to closed-canopy forest is more effective than that which proceeds more slowly.

Twelve of the papers treat bottomland hardwood or other lowland forest types. Among these, Ribbeck and Hunter

(1994) note that many bird species of highest conservation priority in the MAV are late successional species; therefore, rapid afforestation will benefit them.

Wesley and others (1976) studied winter birds in 44 cottonwood plantations aged 4, 5, or 6 years old, within primarily forested landscapes in the MAV. In an associated study, Tomlinson (1977) conducted WBPS on five plots, including three on two cottonwood plantations and two mature hardwood controls. Six species preferred the cottonwood plantations, 4 showed no preference, and 15 preferred the natural stands (Tomlinson 1977, Wesley and others 1976). Wesley and others (1976) compared bird communities in plantations to those in nearby mature natural stands, but not in plowed or harvested agricultural fields. The winter bird community of agricultural fields in the MAV is a simple one (table 2).

In 1994, Twedt conducted three WBPS in planted cottonwoods surrounded by agricultural fields on Fidler Managed Plantation, near Fidler, MS.³ He found 35 species on the plots, 4 of which were not found by Tomlinson (1977) or Wesley and others (1976): red-shouldered hawk, northern bobwhite, chipping sparrow, and Lincoln's sparrow.

Litvaitis (1993) and Sepik and Blumenstock (1993) note that as landscapes change from primarily agricultural to primarily forested settings, species of early successional vegetation benefit in the short term and ultimately decline to the point that specific manipulative action is required to maintain their populations on the landscape. This test does not refute the hypothesis that early successional species will benefit less from afforestation than later successional species.

Empirical Observations

Survey—Table 2 lists the winter bird community Twedt observed in agricultural fields in the MAV. In his surveys of early successional habitats, Broerman tracked occurrence of eight species of birds (table 3). These species are relatively rare in the MAV and of specific conservation interest. Two species—sedge wren and Le Conte's sparrow—occurred on more than half of the surveyed sites (table 3). Each species is associated with grasslands or with herbaceous vegetation in the earliest stages of forest succession (Hamel 1992); neither appears in the cottonwood plantations studied by Wesley and others (1976), Tomlinson (1977), or Twedt.⁴

Experimental test-species occurrence—Vegetation structure on the plots at Sharkey Site differs by treatment at 5 years after establishment. On the NUR treatments, cottonwood trees approach 10 m or more in height. Nuttall oak seedlings are approximately 3 to 4 m tall in the PLN and 1 to 3 m tall in the SOW. On the NAT, few woody stems exceed the 1- to 3-m height of the herbaceous vegetation. These differences in structure are consistent with the

Table 2—Bird species commonly found in fallow agricultural fields in the Mississippi Delta in the winter from three Winter Bird Populations Studies conducted in 1994^a

Common and scientific names	Bird density ^b	
	Per km ²	
Great blue heron <i>Ardea herodias</i> Linn.	0.1 ±	0.1
Northern harrier <i>Circus cyaneus</i> (L.)	.4 ±	.2
Cooper's hawk <i>Accipiter cooperii</i> (Bonaparte)	.4 ±	.4
Red-tailed hawk <i>Buteo jamaicensis</i> (Gmelin)	.7 ±	.3
American kestrel <i>Falco sparverius</i> L.	.1 ±	.1
Killdeer <i>Charadrius vociferus</i> L.	23.6 ±	21.0
Common snipe <i>Gallinago gallinago</i> (L.)	.1 ±	.1
Rock dove <i>Columba livia</i> Gmelin	9.7 ±	9.7
Mourning dove <i>Zenaida macroura</i> (L.)	17.1 ±	11.1
Red-bellied woodpecker <i>Melanerpes carolinus</i> (L.)	.2 ±	.1
Loggerhead shrike <i>Lanius ludovicianus</i> L.	1.9 ±	1.3
Blue jay <i>Cyanocitta cristata</i> (L.)	.9 ±	.7
Horned lark <i>Eremophila alpestris</i> (L.)	17.3 ±	12.1
European starling <i>Sturnus vulgaris</i> L.	189.0 ±	189.0
Vesper sparrow <i>Pooecetes gramineus</i> (Gmelin)	.4 ±	.4
Savannah sparrow <i>Passerculus sandwichensis</i> (Gmelin)	61.9 ±	57.7
Song sparrow <i>Melospiza melodia</i> (Wilson)	3.8 ±	2.7
White-throated sparrow <i>Zonotrichia albicollis</i> (Gmelin)	10.2 ±	9.5
White-crowned sparrow <i>Z. leucophrys</i> (Forster)	.7 ±	.7
Dark-eyed junco <i>Junco hyemalis</i> (L.)	.1 ±	.1
Northern cardinal <i>Cardinalis cardinalis</i> (L.)	8.2 ±	7.1
Red-winged blackbird <i>Agelaius phoeniceus</i> (L.)	2.3 ±	1.4
Eastern meadowlark <i>Sturnella magna</i> (L.)	1.2 ±	1.2
Mean density	350.4 ±	210.9
Mean species richness	13.3 ±	2.8

³ Twedt, Daniel. 1999. Unpublished data. On file with: U.S. Department of the Interior, Geological Survey, Patuxent Wildlife Research Center, 2524 South Frontage Road, Vicksburg, MS.

⁴ Twedt, Daniel. 1994. Unpublished data. On file with: U.S. Department of the Interior, Geological Survey, Patuxent Wildlife Research Center, 2524 South Frontage Road, Vicksburg, MS.

^a Twedt, Daniel. 1994. Unpublished field notes. On file with: U.S. Department of Interior, Geological Survey, Patuxent Wildlife Research Center, 2524 South Frontage Road, Vicksburg, MS.

^b Plus or minus standard error.

Table 3—Bird species observed on 273 trips to 59 afforestation sites in 9 counties in Mississippi from October 17, 1996 to April 24, 2000, by F. Broerman

Common and scientific names ^a	Sites	Birds/trip
	- - - - Number - - - -	
American bittern <i>Botaurus lentiginosus</i> (Rackett)	14	2.3
Yellow rail <i>Coturnicops noveboracensis</i> (Gmelin)	4	1.0
Sora <i>Porzana carolina</i> (L.)	16	5.1
Short-eared owl <i>Asio flammeus</i> (Pontoppidan)	21	5.6
Sedge wren <i>Cistothorus platensis</i> (Latham)	33	2.5
Marsh wren <i>C. palustris</i> (Wilson)	20	2.4
Field sparrow <i>Spizella pusilla</i> (Wilson)	10	3.1
Le Conte's sparrow <i>Ammodramus leconteii</i> (Audubon)	36	4.5

^a In addition to the species listed, Broerman did not count individuals of savannah, swamp, and song sparrows, or of eastern meadowlarks, all of which were numerous on the surveyed sites.

intensity of the management of the plots at establishment, with the age of the propagules when planted, and with the growth rates of the different species planted. Vegetation structure has accumulated more rapidly in plots in which more intense effort was made to establish woody vegetation.

Hamel found a total of 51 bird species in the 1999 to 2000 WBPS at the Sharkey Site (table 4). We examine the occurrence of individual species as a function of their association with vegetation structure, as well as with respect to their conservation priority. The ANOVA of species richness among treatments revealed that significantly more species (30 plus or minus 4.6 S.D. vs. 11.7 plus or minus 1.8 S.D.) occurred in the NUR treatment than in the others ($F = 54.2$, d.f. = 5,6, $P < 0.0001$ (table 4). The ANOVA of total abundance was similar among treatments ($F = 1.13$, d.f. = 5,6, $P < 0.43$ (table 4). Thus 5 years after establishment, the treatment with the greatest development of vegetation structure (NUR) harbored the greatest number of bird species, although it did not harbor a greater number of individuals than the other treatments. Among the 28 species found only in the NUR treatment, 12 foraged on trees (Hamel 1992). Four of the five species never found in the NUR foraged on the ground or herbaceous vegetation. These results are consistent with the hypothesis that bird species occurrence reflects physical vegetation structure.

As vegetation structure develops, avian species appear that are associated with that structure, such as Eastern phoebe and yellow-rumped warbler. These birds, which forage from

trees or at variable heights, were recorded only in the NUR plots in 1998 to 1999. In 1999 to 2000, these birds occurred rarely in the emerging woody vegetation of the other treatments, particularly the PLN. In both years, species associated primarily with open vegetation also occurred beneath the trees in the NUR, notably song sparrow, swamp sparrow, and red-winged blackbird.

Other species associated with open habitats, such as Northern harrier, sedge wren, savannah sparrow, Le Conte's sparrow, and eastern meadowlark rarely occurred in the NUR treatment. For example, sedge wren was found regularly in all open habitats (54 of 72 visits), but in the NUR only once (of 24 visits) in a very open spot near the edge where flooding had killed some of the planted cottonwoods. An association between occurrence and physical vegetation structure does not explain why Le Conte's sparrow occurred primarily in one spot overlapping the border of a NAT and a PLN plot. The distribution of this bird does not seem to be a straightforward response to the afforestation treatments in this experiment.

Experimental test-species conservation priority—The Partners in Flight (PIF) offers a generally accepted system for recording bird species conservation priority (Partners in Flight 2000). The PIF system gives each species a priority ranking score based on several aspects of occurrence, abundance, and threats to its population. These priority scores are recorded for physiographic areas in which a species breeds, but not for areas where it only winters (Colorado Bird Observatory 1999); the maximum possible score is 35 the minimum is 7 (Carter and others 2000). When possible, we used the PIF conservation priority scores from the MAV for this analysis; where scores were unavailable for the MAV, we used the score from a representative physiographic region in the breeding range of the species. Nuttle and others (2000) have used PIF concern scores similarly.

While bird species richness increases with vegetation structure in the Sharkey Site, conservation priority of individual species does not (table 4). Using the PIF priority rankings, several observations are suggestive. First, the average conservation priority of all 51 species recorded on the WBPS is 15.2. The average ranking of all species found in common among all treatments is a similar 15.5. Second, 28 species that were unique to the NUR treatment average 14.6, a slightly lower value. Third, 5 species not found in the NUR average 16.6, a value higher than the average. When the sedge wren is included in this group, average priority ranking of species found in the other treatments increases to 17.3. Fourth, among the six species of highest conservation priority, two never occurred in the NUR and one occurred only in the NUR. Therefore, rapid afforestation provides winter habitat for a number of species quickly, at the expense of a few high-priority species found in early successional habitats.

CONCLUSIONS

Extensive restoration of forests in the MAV may provide demonstrable, albeit unintended, benefits to birds that winter within afforested sites in early successional stages. This data set illustrates that bird species composition in the MAV

Table 4—Foraging site, abundance on different afforestation treatments, and conservation priority of bird species recorded on four afforestation treatments during winter 1999 to 2000 at the Sharkey Large-Scale Demonstration Project, Sharkey County, Mississippi

Common and scientific names	Foraging site ^e	Abundance in treatment ^d				Conservation priority rating ^f
		NAT	SOW	PLN	NUR	
Northern harrier <i>Circus cyaneus</i> (L.)	G, H	13	13	13	1	21 ^{aa}
Red-tailed hawk <i>Buteo jamaicensis</i> (Gmelin)	G, H	2.1	9.9	2.6	2.1	12
American kestrel <i>Falco sparverius</i> L.	G	.5	0	0	.5	12
Northern bobwhite <i>Colinus virginianus</i> (L.)	G	1	0	0	0	20
Common snipe <i>Gallinago gallinago</i> (L.)	W	2.1	1	.5	0	13 ^{aa}
American woodcock <i>Scolopax minor</i> Gmelin	G	0	0	.5	.5	19
Mourning dove <i>Zenaida macroura</i> (L.)	G	0	0	0	2.6	14
Great horned owl <i>Bubo virginianus</i> (Gmelin)	G	0	0	0	.5	12
Barred owl <i>Strix varia</i> Barton	V	0	0	0	2.6	16
Red-bellied woodpecker ^a <i>Melanerpes carolinus</i> (L.)	T	0	0	0	2.1	17
Yellow-bellied sapsucker ^a <i>Sphyrapicus varius</i> (L.)	T	0	0	0	1	16 ^{aa}
Downy woodpecker ^a <i>Picoides pubescens</i> (L.)	T, B	0	0	0	14.6	14
Hairy woodpecker ^a <i>P. villosus</i> (L.)	T	0	0	0	1.6	14
Northern flicker ^a <i>Colaptes auratus</i> (L.)	G, T	0	0	0	7.8	16
Eastern phoebe <i>Sayornis phoebe</i> (Latham)	V	.5	0	2.1	19.3	15
Loggerhead shrike <i>Lanius ludovicianus</i> L.	G	4.7	3.1	8.3	.5	19
Blue jay <i>Cyanocitta cristata</i> (L.)	T	0	0	0	2.1	13
American crow <i>Corvus brachyrhynchos</i> Brehm	G	0	0	0	.5	10
Carolina chickadee ^a <i>Poecile carolinensis</i> (Audubon)	T	0	0	0	13	20
Tufted titmouse ^b <i>Baeolophus bicolor</i> (L.)	T	0	0	0	.5	14
Carolina wren ^b <i>Thryothorus ludovicianus</i> (Latham)	G, B	0	0	0	9.9	17
Winter wren ^b <i>Troglodytes troglodytes</i> (L.)	G	0	.5	0	8.8	14 ^{bb}
Sedge wren <i>Cistothorus platensis</i> (Latham)	H	23.4	23.7	21.4	.5	21
Golden-crowned kinglet <i>Regulus satrapa</i> Lichtenstein	T	0	0	0	3.1	17 ^{bb}

continued

Table 4—Foraging site, abundance on different afforestation treatments, and conservation priority of bird species recorded on four afforestation treatments during winter 1999 to 2000 at the Sharkey Large-Scale Demonstration Project, Sharkey County, Mississippi (continued)

Common and scientific names	Foraging site ^e	Abundance in treatment ^d				Conservation priority rating ^f
		NAT	SOW	PLN	NUR	
Ruby-crowned kinglet ^a						
<i>Regulus calendula</i> (L.)	T, B	0	0	0	7.3	16 ^{bb}
Eastern bluebird ^a						
<i>Sialia sialis</i> (L.)	G	0	0	0	2.1	14
Hermit thrush ^a						
<i>Catharus guttatus</i> (Pallas)	G, B	0	0	0	3.1	16 ^{bb}
American robin ^a						
<i>Turdus migratorius</i>	G, B, T	0	0	0	13	9
Northern mockingbird						
<i>Mimus polyglottos</i> (L.)	G, B	0	0	.5	0	14
Brown thrasher						
<i>Toxostoma rufum</i> (L.)	G, B	0	0	0	.5	17
Cedar waxwing						
<i>Bombycilla cedrorum</i> Vieillot	B, T	0	0	0	2.1	12 ^{aa}
Yellow-rumped warbler ^a						
<i>Dendroica coronata</i> (L.)	V, T	0	0	.5	65.6	16 ^{bb}
Palm warbler						
<i>Dendroica palmarum</i> (Gmelin)	G, H, B	0	0	0	.5	16 ^{cc}
Common yellowthroat						
<i>Geothlypis trichas</i> (L.)	H, B	.5	2.6	1	.5	16
Eastern towhee						
<i>Pipilo erythrophthalmus</i> (L.)	G, B	0	0	0	.5	15
Field sparrow						
<i>Spizella pusilla</i> (Wilson)	H	0	2.1	0	.5	20
Savannah sparrow						
<i>Passerculus sandwichensis</i> (Gmelin)	G, H	94.8	74.5	140.1	0	13 ^{aa}
Le Conte's sparrow						
<i>Ammodramus leconteii</i> (Audubon)	H	1.6	0	4.7	0	23 ^{aa}
Fox sparrow						
<i>Passerella iliaca</i> (Merrem)	G	0	.5	.5	15.1	12 ^{bb}
Song sparrow ^c						
<i>Melospiza melodia</i> (Wilson)	G, H	68.2	78.1	53.1	30.2	12
Swamp sparrow ^c						
<i>M. georgiana</i> (Latham)	H	78.6	86.4	124	55.2	16 ^{aa}
White-throated sparrow ^c						
<i>Zonotrichia albicollis</i> (Gmelin)	G, B	0	0	0	43.2	14 ^{cc}
White-crowned sparrow						
<i>Z. leucophrys</i> (Forster)	G, H	0	0	0	2.1	12 ^{bb}
Dark-eyed junco ^c						
<i>Junco hyemalis</i> (L.)	G	0	0	0	10.4	14 ^{cc}
Northern cardinal						
<i>Cardinalis cardinalis</i> (L.)	G, B	0	0	0	5.2	12
Red-winged blackbird ^c						
<i>Agelaius phoeniceus</i> (L.)	G, H	241.7	18.2	63.5	252.6	12
Eastern meadowlark						
<i>Sturnella magna</i> (L.)	G, H	50	47.9	61.5	4.2	17

continued

Table 4—Foraging site, abundance on different afforestation treatments, and conservation priority of bird species recorded on four afforestation treatments during winter 1999 to 2000 at the Sharkey Large-Scale Demonstration Project, Sharkey County, Mississippi (continued)

Common and scientific names	Foraging site ^e	Abundance in treatment ^d				Conservation priority rating ^f
		NAT	SOW	PLN	NUR	
Rusty blackbird						
<i>Euphagus carolinus</i> (Müller)	G	0	0	0	8.3	16 ^{cc}
Brewer's blackbird						
<i>E. cyanocephalus</i> (Wagler)	G	0	0	0	1.6	15 ^{aa}
Common grackle						
<i>Quiscalus quiscula</i> (L.)	G	0	0	.5	222.4	16
American goldfinch ^c						
<i>Carduelis tristis</i> (L.)	V	3.6	0	3.1	16.7	12
Unknown species	NA	6.3	5.2	5.2	7.3	NA
Mean total density per treatment	NA	592.6	366.7	507.2	865.3	NA
Mean no. of species per treatment	NA	11	11	13	30	NA
Total species per treatment	NA	17	14	19	47	51
Unique species per treatment	NA	1	0	1	28	NA

NA = not applicable; NAT = natural succession and regeneration control; SOW = sown Nuttall oak acorns, 2500 per hectare; PLN = planted Nuttall oak seedlings, 750 per hectare; NUR = planted cottonwood stem cuttings, 750 per hectare, followed by underplanting of Nuttall oak seedlings, 375 per hectare, 2 years later.

^a Species associated with cottonwood plantations, which are listed by Wesley and others (1976) as preferring natural stands.

^b Species associated with cottonwood plantations, which are listed by Wesley and others (1976) as preferring neither plantations nor natural stands.

^c Species listed by Wesley and others (1976) as preferring cottonwood plantations.

^d Entries reflect mean abundance as birds per km². Boldfaced numbers indicate mean abundance different from other treatments by one-way analysis of variance with treatment as main effect, adjusted for an experiment-wide error rate of 0.10 using sequential Bonferroni correction. Standard error values available on request from Hamel (lead author of publication).

^e Foraging sites listed as B = bush, G = ground, H = herbaceous vegetation, T = tree, V = various heights, W = water (Hamel 1992).

^f Conservation priority ratings are the Partners in Flight Concern Scores (Colorado Bird Observatory 1999) for the Mississippi Alluvial Valley, with the exception of ^{aa} = Drift Prairie, ^{bb} = Central Rocky Mountains, and ^{cc} = Great Lakes Transition physiographic regions.

follows vegetation structure and development rate in afforestation of abandoned agricultural lands. However, as woody vegetation develops, some bird species of herbaceous vegetation disappear. Perhaps more importantly, the early successional avian species that specialize on herbaceous vegetation are of higher than average conservation priority among the birds found in afforestation areas.

Different species of mammals and birds respond positively to the structure available at different times during succession. Thus, managers must decide on the species and communities they wish to favor. Winter avian communities of early successional stages have been considered heretofore only in passing. However, because birds occur in areas where vegetation structure is dominated by nonagricultural herbaceous plants, they include species otherwise rare or absent from the MAV landscape. To provide bird habitat through the full successional sere, proactive managers may wish to dedicate certain areas to natural succession or to maintenance of herbaceous vegetation.

Forest land managers and biodiversity preservationists make clear the essential relationship between management practices and management objectives (McCullough 1994, Noss 1999). Management (1) is a conscious, goal-directed activity with goals specified in advance, and (2) employs practices to achieve goals within an acceptable period of time. Managers can apply a variety of afforestation methods under different conditions to achieve different management objectives. Managerial discretion influences the rate of structural development of afforestation efforts. Inclusion of management goals to produce habitats in afforestation for early successional species is certainly possible, as is inclusion of management goals to maximize the rate of development of forest vegetation structure and the birds associated with it.

ACKNOWLEDGMENTS

This work has benefited from the generous field assistance of the staff and residents of Sunflower Landing, Dublin, MS. Mark Woodrey and Jeff Buler of the Mississippi Museum of

Natural Science aided in the fieldwork, as did a number of other collaborators on the Sharkey Large-Scale Restoration Project.

LITERATURE CITED

- Anderson, R.C.; Katz, A.J. 1993. Recovery of browse-sensitive tree species following release from white-tailed deer *Odocoileus virginianus* Zimmerman browsing pressure. *Biological Conservation*. 63(3): 203–208.
- Askins, R.A.; Philbrick, M.J. 1987. Effect of changes in regional forest abundance on the decline and recovery of a forest bird community. *Wilson Bulletin*. 99(1): 7–21.
- Bauch, J.; Dunisch, O.; Gasparotto, L., eds. 1999. Investigations on tree species suitable for the recultivation of degraded land areas in central Amazonia: Brazilian-German cooperation within the project 'studies on human impacts on forests and floodplains in the tropics' (SHIFT) supported by the Ministry of Education, Science, Research, and Technology, Bonn, FRG (0339638) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brasília, Brazil (ENV 42) and Instituto Brasileiro do Meio Ambiente e Recursos Naturais Renováveis (IBAMA). No. 193. Hamburg, Germany: Mitteilungen der Bundesforschungsanstalt für Forst- und Holzwirtschaft. 138 p.
- Bielefeldt, J.; Rosenfield, R.N. 1994. Summer birds of conifer plantations in southeastern Wisconsin. *Passenger Pigeon*. 56(2): 123–135.
- Boyle, J.R. 1999. Planted forests: views and viewpoints. *New Forests*. 17(1/3): 5–9.
- Burgess, P.J. 1999. Effects of agroforestry on farm biodiversity in the UK. *Scottish Forestry*. 53(1): 24–27.
- Cannell, M.G.R. 1999a. Environmental impacts of forest monocultures: water use, acidification, wildlife conservation, and carbon storage. *New Forests*. 17(1/3): 239–262.
- Cannell, M.G.R. 1999b. Forests, Kyoto and climate. *Outlook on agriculture*. 28(3): 171–177.
- Cantrell, M.A.; Britcher, J.J.; Hoffman, E.L. 1995. Red-cockaded woodpecker management initiatives at Fort Bragg Military Installation. In: Kulhavy, D.L.; Hooper, R.G.; Costa, R., eds. Red-cockaded woodpecker: recovery, ecology and management. Red-cockaded woodpecker symposium III: species recovery, ecology and management; 1993 January 24–28; North Charleston, SC. Nacogdoches, TX: Stephen F. Austin State University, Center for Applied Studies, College of Forestry: 89–97.
- Carter, M.F.; Hunter, W.C.; Pashley, D.N.; Rosenberg, K.V. 2000. Setting conservation priorities for landbirds in the United States: the partners in flight approach. *Auk*. 117(2): 541–548.
- Chang, ChingCheng. 1999. Carbon sequestration cost by afforestation in Taiwan. *Environmental Economics and Policy Studies*. 2(3): 199–213.
- Colorado Bird Observatory. 2000. Colorado Bird Observatory PIF database. Barr Lake State Park, CO: Colorado Bird Observatory. [Available on the Web: <http://www.cbobirds.org/pif/index.html>].
- Conner, R.N.; Rudolph, D.C. 1995. Losses of red-cockaded woodpecker cavity trees to southern pine beetles. *Wilson Bulletin*. 107(1): 81–92.
- Cronquist, M.J.; Brooks, R.P. 1991. Use of avian and mammalian guilds as indicators of cumulative impacts in riparian-wetland areas. *Environmental Management*. 15(5): 701–714.
- DeGraaf, R.M. 1987. Managing northern hardwoods for breeding birds. In: Nyland, R.D., ed. Proceedings of the silvicultural symposium managing northern hardwoods; 1986 June 23–25; Syracuse, NY. Misc. Publ. 13 (ESF 87–002). Syracuse, NY: Syracuse University, Faculty of Forestry: 348–362.
- DeGraaf, R.M.; Yamasaki, M.; Leak, W.B.; Lanier, J.B. 1992. New England wildlife: management of forested habitats. Gen. Tech. Rep. NE–144. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 271 p.
- Dickson, J.G., ed. 1992. The wild turkey: biology and management. Harrisburg, PA: Stackpole Books. 463 p.
- Fitzgerald, S.M.; Tanner, G.W. 1992. Avian community response to fire and mechanical shrub control in south Florida. *Journal of Range Management*. 45(4): 396–400.
- Flebbe, P.A.; Dolloff, C.A. 1995. Trout use of woody debris and habitat in Appalachian wilderness streams of North Carolina. *North American Journal of Fisheries Management*. 15: 579–590.
- Gaines, G.D.; Franzreb, K.E.; Allen, D.H. [and others]. 1995. Red-cockaded woodpecker management on the Savannah River Site: a management/research success story. In: Kulhavy, D.L.; Hooper, R.G.; Costa, R., eds. Red-cockaded woodpecker: recovery, ecology and management. Red-cockaded woodpecker symposium III: species recovery, ecology and management; 1993 January 24–28; North Charleston, SC. Nacogdoches, TX: Stephen F. Austin State University, Center for Applied Studies, College of Forestry: 81–88.
- Hamel, P.B. 1992. A land manager's guide to the birds of the South. Atlanta: U.S. Department of Agriculture, Forest Service, Southern Region; Chapel Hill, NC: The Nature Conservancy, Southeastern Region. 367 p.
- Harrington, C.A. 1999. Forests planted for ecosystem restoration or conservation. *New Forests*. 17(1/3): 175–190.
- Harris, L.D.; Scheck, J. 1991. From implications to applications: the dispersal corridor principle applied to the conservation of biological diversity. In: Saunders, D.A.; Hobbs, R.J., eds. *Nature conservation 2: the role of corridors*. Chipping Norton, Australia: Surrey Beatty & Sons: 189–220.
- Helmer, W. 1999. [Establishment of broadleaved floodplain forest]. *Opkomst van het hardhoutooibos*. *Nederlands Bosbouw tijdschrift*. 71(3): 130–134.
- Houston, A.E. 1991. Beaver control and reforestation of drained beaver impoundments. Knoxville, TN: University of Tennessee. 143 p. Ph.D. dissertation.
- Hurst, G.A.; Bourland, T.R. 1996. Breeding birds on bottomland hardwood regeneration areas on the Delta National Forest. *Journal of Field Ornithology*. 67: 181–187.
- James, F.C. 1971. Ordinations of habitat relationships among breeding birds. *Wilson Bulletin*. 83: 215–236.
- Joslin, J.D.; Schoenholtz, S.H. 1997. Measuring the environmental effects of converting cropland to short-rotation woody crops: a research approach. *Biomass and Bioenergy*. 13(4/5): 301–311.
- Kaminski, R.M.; Alexander, R.W.; Leopold, B.D. 1993. Wood duck and mallard winter microhabitats in Mississippi hardwood bottomlands. *Journal of Wildlife Management*. 57(3): 562–570.

- Kolb, H. 1965. The Audubon winter bird population study. *Audubon Field Notes*. 19(3): 432–434.
- Litvaitis, J.A. 1993. Response of early successional vertebrates to historic changes in land use. *Conservation Biology*. 7(4): 866–873.
- McCullough, D.R. 1994. Importance of population data in forest management planning. *Forestry Chronicle*. 70(5): 533–537.
- Mitchell, L.J. 1989. Effects of clearcutting and reforestation on breeding bird communities of baldcypress-tupelo wetlands. Raleigh, NC: North Carolina State University. 92 p. M.S. thesis.
- Mladenoff, D.J.; White, M.A.; Crow, T.R.; Pastor, J. 1994. Applying principles of landscape design and management to integrate old-growth forest enhancement and commodity use. *Conservation Biology*. 8(3): 752–762.
- Mladenoff, D.J.; White, M.A.; Pastor, J.; Crow, T.R. 1993. Comparing spatial pattern in unaltered old-growth and disturbed forest landscapes. *Ecological Applications*. 3(2): 294–306.
- Mueller, A.J.; Loesch, C.R.; Twedt, D.J. 1995. Development of management objectives for breeding birds in the Mississippi Alluvial Valley. In: Bonney, R.; Pashley, D.N.; Cooper, R.J.; Niles, L., eds. *Proceedings of the Partners in Flight international workshop strategies for bird conservation: the Partners in Flight planning process*; 1995 October 1–5; Cape May, NJ. Ithaca, NY: Cornell Lab of Ornithology. [Various pagination]. [Available on the Web: <http://www.birds.cornell.edu/pifcapemay/mueller.htm>].
- National Council of the Paper Industry for Air and Stream Improvement, Inc. (NCASI). 1993. *Forestry, wildlife, and habitat in the East: an annotated bibliography, 1986–1990*. NCASI Tech. Bull. 651. Research Triangle Park, NC: National Council of the Paper Industry for Air and Stream Improvement, Inc. 340 p.
- National Council of the Paper Industry for Air and Stream Improvement, Inc. (NCASI). 1999. *Forestry, wildlife, and habitat in the East: an annotated bibliography, 1991–1995*. NCASI Tech. Bull. 781. Research Triangle Park, NC: National Council of the Paper Industry for Air and Stream Improvement, Inc. 753 p.
- Niyaz, S.; Nigi, T.; Koshika, K. 1999. [Growth of planted subsidized trees in natural forests in the central Tianshan Mountains and damage caused by grazing cattle]. *Research Bulletin of the Hokkaido University Forests*. 56(1): 94–104.
- Noss, R.F. 1999. Assessing and monitoring forest biodiversity: a suggested framework and indicators. *Forest Ecology and Management*. 115(2/3): 135–146.
- Nuttle, T.J. 1997. Response of breeding bird communities to afforestation of hardwood bottomland sites in Mississippi. Mississippi State, MS: Mississippi State University. 68 p. M.S. thesis.
- Nuttle, T.J.; Burger, L.W. 1995. Breeding bird response to hardwood bottomland restoration in the Mississippi Alluvial Valley. [Abstract]. In: Program of Wildlife Society second annual conference: excellence in wildlife stewardship through science and education; 1995 September 12–17; Portland, OR: 92–93.
- Nuttle, T.; Leidolf, A.; Burger, L.W. 2000. Using Partners in Flight species concern scores to compare the conservation value of bird communities [Abstract]. In: One hundred eighteenth stated meeting of the American Ornithologists' Union; 2000 August 14–19; St. John's, Newfoundland, Canada. St. John's, Newfoundland, Canada: American Ornithologists' Union: 26.
- O'Brien-White, S.; Thomason, C.S. 1995. Evaluating fish habitat in a South Carolina watershed using GIS. *Proceedings of the annual conference of the Southeastern Association of Fish and Wildlife Agencies*; 1995 September 23–27; Nashville, TN. Tallahassee, FL: Southeastern Association of Fish and Wildlife Agencies. 49: 153–166.
- Pande, V.C.; Nambiar, K.T.N.; Singh, H.B. 1999. Afforestation on community land in the semi-arid tropics of Gujarat—an economic analysis. *Indian Forester*. 125(2): 212–218.
- Partners in Flight. 2000. Partners in Flight – U.S. home page. Patuxent National Wildlife Refuge, MD: U.S. Department of the Interior, Biological Resources Division. [Various pagination]. [Available on the Web: <http://www.partnersinflight.org>].
- Provencher, L.; Demarest, D.; Brennan, L. [and others]. 1995. Avian community responses to herbaceous invertebrates and vegetation in burned old-growth and fire-suppressed longleaf pine habitats: experimental design and pretreatment results [Abstract]. In: Program of Wildlife Society second annual conference: excellence in wildlife stewardship through science and education; 1995 September 12–17; Portland, OR. Portland, OR: The Wildlife Society: 105.
- Reinecke, K.J. 1994. Regional waterfowl habitat trends and implications for neotropical migratory birds. In: Smith, W.P.; Pashley, D.N., comps., eds. *A workshop to resolve conflicts in the conservation of migratory landbirds in bottomland hardwood forests*; 1993 August 9–10; Tallulah, LA. Gen. Tech. Rep. SO–114. New Orleans: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 20–24.
- Ribbeck, K.F.; Hunter, W.C. 1994. Is reforestation an adequate restoration of bottomland hardwood functions for the needs of neotropical migratory birds? In: Smith, W.P.; Pashley, D.N., comps., eds. *A workshop to resolve conflicts in the conservation of migratory landbirds in bottomland hardwood forests*; 1993 August 9–10; Tallulah, LA. Gen. Tech. Rep. SO–114. New Orleans: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 25–26.
- Robinson, G.R.; Handel, S.N. 1993. Forest restoration on a closed landfill: rapid addition of new species by bird dispersal. *Conservation Biology*. 7(2): 271–278.
- Root, K.V.; Breininger, D.R.; Bowman, R.; Swain, H.M. 1995. What can population viability analysis (PVA) tell us about the probable success of a spatially explicit reserve design: the Florida scrub jay in Brevard County as a case study [Abstract]. *Bulletin of the Ecological Society of America*. 76(2 Suppl. Part 2): 229.
- SAS Institute Inc. 1989. *SAS/STAT user's guide*. Version 6. 4th ed. Cary, NC: SAS Institute Inc. 846 p. Vol. 2.
- Schmalzer, P.A. 1993. Characteristics of long unburned stands of oak-saw palmetto scrub [Abstract]. *Bulletin of the Ecological Society of America*. 74(2 Suppl.): 428.
- Schmalzer, P.A. 1994. Restoring unburned shrublands: an example using cutting and burning in oak-saw palmetto scrub [Abstract]. *Bulletin of the Ecological Society of America*. 75(2 Suppl.): 206.
- Schmalzer, P.A.; Breininger, D.R.; Adrian, F.W. [and others]. 1994. Development and implementation of a scrub habitat compensation plan for Kennedy Space Center. Tech. Memo. 109202. Kennedy Space Center, FL: National Aeronautics and Space Administration. 54 p.
- Scholtens, L.J.R. 1998. Environmental, developmental and financial risks of tropical timber plantation investment funds. *Natural Resources Forum*. 22(4): 271–277.

- Schweitzer, C.J.; Stanturf, J.A.; Shepard, J.P. [and others]. 1997. Large-scale comparison of reforestation techniques commonly used in the Lower Mississippi Alluvial Valley: first year results. In: Pallardy, S.G.; Cecich, R.A.; Garrett, H.; Johnson, P.S., eds. Proceedings of the 11th central hardwood forest conference; 1997 March 23–26; Columbia, MO. Gen. Tech. Rep. NC–188. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station: 313–320.
- Sepik, G.F.; Blumenstock, B. 1993. Management and research on the American woodcock at the Moosehorn National Wildlife Refuge [Abstract]. In: Longcore, J.R.; Sepik, G.F., eds. Proceedings of the eighth American woodcock symposium; 1993; West Lafayette, IN. Biol. Rep. 16. Washington, DC: U.S. Department of the Interior, Fish and Wildlife Service, National Biological Survey: 136–137.
- Simberloff, D. 1993. Effects of fragmentation on some Florida ecosystems, and how to redress them. In: Saunders, D.A.; Hobbs, R.J.; Ehrlich, P.R., eds. Nature conservation 3: reconstruction of fragmented ecosystems. Chipping Norton, Australia: Surrey Beatty & Sons: 179–187.
- Sweeney, B.W. 1993. Effects of streamside vegetation on macroinvertebrate communities of White Clay Creek in Eastern North America. Proceedings of the Academy of Natural Sciences of Philadelphia. 144: 291–340.
- Vora, R.S. 1994. Integrating old-growth forest into managed landscapes: a northern Great Lakes perspective. *Natural Areas Journal*. 14(2): 113–123.
- Watson, J.C.; Hooper, R.G.; Carlson, D.L. [and others]. 1995. Restoration of the red-cockaded woodpecker population on the Francis Marion National Forest: three years post Hugo. In: Kulhavy, D.L.; Hooper, R.G.; Costa, R., eds. Red-cockaded woodpecker: recovery, ecology and management. Red-cockaded woodpecker symposium III: species recovery, ecology and management; 1993 January 24–28; North Charleston, SC. Nacogdoches, TX: Stephen F. Austin State University, Center for Applied Studies, College of Forestry: 172–182.
- Weaver, K.M.; Pelton, M.R. 1994. Denning ecology of black bears in the Tensas River basin of Louisiana. *International Conference on Bear Research and Management*. 9(1): 427–433.
- Weaver, K.M.; Tabberer, D.K.; Moore, L.U., Jr. [and others]. 1990. Bottomland hardwood forest management for black bears in Louisiana. Proceedings of the annual conference of the Southeastern Association of Fish and Wildlife Agencies; 1990 October 21–24; Richmond, VA. Tallahassee, FL: Southeastern Association of Fish and Wildlife Agencies. 44: 342–350.
- Wesley, D.E.; Perkins, C.J.; Sullivan, D.A. 1976. Preliminary observations of cottonwood plantations as wildlife habitat. In: Thielges, B.A.; Land, S.B., Jr. eds. Proceedings of the symposium on eastern cottonwood and related species; 1976 September 28–October 2; Greenville, MS. New Orleans: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 460–476.
- Willoughby, I.; McDonald, H.G. 1999. Vegetation management in farm forestry: a comparison of alternative methods of inter-row management. *Forestry (Oxford)*. 72(2): 109–121.
- Wilson, C.W.; Masters, R.E.; Buehner, G.A. 1995. Breeding bird response to pine-grassland community restoration for red-cockaded woodpeckers. *Journal of Wildlife Management*. 59(1): 56–67.